



Pink

THE PINK-AGING TREATMENT

PINK-AGING

Brightening · Anti-Pigmentation · Anti-Aging · Replenishing Skin barrier

The **Pink-Aging treatment** is a dermatological solution for professionals that stimulates the vitalization of fibroblast cells, which are the basic structural frameworks of tissues, by improving skin metabolism via a dynamic activation of its 56 components.



Pink Main Components


The Pink-Aging treatment is a dermatological solution composed of 56 components that helps improve the vitalization fibroblast cells, which are the basic structural frameworks of tissues.



Pink Special Features

Skin Tone up

- ▶ Contains **glutathione**, which brightens skin and removes free radicals ¹⁷
- ▶ Contains **3 kinds of enzyme**, which contribute to the synthesis of glutathione in skin ¹⁸
- ▶ Contains **10 kinds of vitamin B complex (vitamin B12)**, which improve skin metabolism ²³



Ref: International Journal of molecular Sciences [2018], NO.9 pp2845-0

Skin Rejuvenation

- ▶ Contains **growth factors, adenosine A2 receptors**, which help quickly recover skin regeneration cycle ^{1,2,6,7,9,10,18,19,20,21}
- ▶ Contains **6 kinds of peptide (EGF, FGF, IGF, TGF- β , cooper tripeptide)**, which help vitalize fibroblast cells, which are the basic structural frameworks of tissues ^{12,13,14}
- ▶ Contains **neuropeptide complex**, which is a growth factor that helps vitalize fibroblast cells and growth of blood vessels, which are essential for a wound healing process

Wound healing effects of neurovax




Ref: BioFD&C R&D Center

Skin Moisturization

- ▶ Contains **amino acid complex**, which increases the content of NMF (natural moisturizing factor) ¹⁸
- ▶ Contains **1.4 million Dalton of high molecule HA**, which can improve skin moisture deficit

Change of skin moisture over time




Ref: Sk Bioland

Skin Energy Activation


- ▶ Contains **vitamin B and amino acid** for coenzyme synthesis
- ▶ Contains **ubiquinone (coenzyme Q10)**, which is the connection for the production of ATP ²³

The process in which coenzyme Q10 affects the electron transport system




1. Masahiro Shiragaki, AL ZOOLOGICAL SCIENCE 17:661-666(2000) 2. Biochem. J. (2006) 400: 225-234 3. Mek, K.K. and Chan, S.Y. (2003) Epidermal growth factor as a biologic switch in hair growth cycle. The journal of biological chemistry, 278(28):26120-26126. 4. Alexandrescu, D. T., Kauffman, C. L., and Dasanu, C. A. (2009) The cutaneous epidermal growth factor network: Can it be translated clinically to stimulate hair growth? Dermatology online journal, 15 (3) 5. McEwee K.J., and Sinclair R. (2008) Hair physiology and its disorders. Drug discovery today, disease mechanisms, 5(2):163-171. 6. American Journal of Physiology: Renal Physiology (2007) 293: F1007-F1017 7. Imamura, T. (2014) Physiological functions and underlying mechanisms of fibroblast growth factor (FGF) family members: recent findings and Implications for their pharmacological application. Biological and pharmaceutical bulletin, 37(7):1081-1089. 8. Stefan Beckett, et al. Surgical Research 139:217-221(2007) 9. The Journal of Biological Chemistry (2001) 276(21): 18185-18192 10. The New England Journal of Medicine (1999) 341(10): 738-746 11. Simeon A, et al. Invest Dermatol. 115(6):962-8(2000) 12. Maquart FX, et al. FEBS Lett 238(2):343-6(1998) 13. Miriam A. al. European Journal of Cell Biology 81: 153-160(2002) 14. Margadant, C. and Sonnenberg, A. (2010) Integrin-TGF- β crosstalk in fibrosis, cancer and wound healing. EMBO Reports, 11(2):97-105. 15. Clinical Cosmetic and Investigational Dermatology 2014;7 267-274 16. Biochim Biophys Acta, 2013 May; 1830(5): 3143-3153. 17. Baden et al. Biochem, Biophysics, Acta 322, p269-278, 1973 18. PDRN improves angiogenesis and wound healing in experimental thermal injury: bio chemical pharmacology 77(2009) 1117-1124 19. PDRN enhance the proliferation of human skin fibroblasts: involvement of A2 purinergic receptor subtypes: Life Science, Vol. 64, No. 18, pp 1661-1674 1999 20. Cell biochemistry and Functio 1999 Life Science 1999 21. Biochim Biophys Acta, 2013 May; 1830(5):3143-53. doi: 10.1016/j.bbagen.2012.09.008. Epub 2012 Sep 17. 35. Journal of the American Academy of Dermatology, 1989, 21권 1호 1p~30p 22. Vitamin B12 Deficiency Induces Imbalance in Melanocytes Homeostasis—A Cellular Basis of Hypocobalaminemia Pigmentary Manifestations. Int J Mol Sci., 2018 Sep 19; 19(9). pii: E2845. doi: 10.3390/ijms19092845. 23. The Effect of Coenzyme Q10 (Ubiquinone) on Random Pattern Skin Flap Survival in Rat Mode—Annals of Plastic Surgery, SCI 2016년 77권, 2호 9p ~ 14p ISSN 0148-7043



Pink Introduction



THE PINK-AGING TREATMENT




RIBESKIN[®]X Multi-needle



- ✓ User-friendly ergonomic design
 - ✓ Easy and accurate release of purposed drug
 - ✓ Enhanced skin permeability in pain-free manner
- *Pink shooter must be used vertically during the application and when being put away. Beware that Pink shooter should not slide away from the skin.

Penetration Rate Comparison



"The use of TURTLEPIN[®]III series (both 0.5mm & 1.0mm) significantly results in better penetration rates of eGFP than in the use of MTS-Roller. [...] Considering the greater convenience and reduced pain experienced in the use of TURTLEPIN[®]III, it may be more suitable for mesotherapy – ideal for both patients and professionals." Korea Research Institute of Bioscience & Biotechnology (KRIBB, 2013), *Comparison study of TURTLEPIN[®]III and mesotherapy-roller

"0.5mm shows higher FL intensity level (30x) and faster penetration speed(20x) in comparison to normal application (NC)." Korea Research Institute of Bioscience & Biotechnology(KRIBB, 2013), *In Vivo&In Vitro – TURTLEPIN[®]III

Pink Treatment Cycles

Phase 1.
Stimulating

3 sessions
with an interval of
1 week to 2 weeks






Phase 2.
Firming

2 sessions
with an interval of
1 month

Phase 3.
Maintaining

1 session
every **3 to 6 months**
regularly

Pink Procedure




Warning

- Avoid exposure to the sunlight. Use of hat, mask, umbrella recommended in outdoors.
- Prior to treatment, avoid use of toner with high contents of alcohol and cream containing AHA, BHA, Vitamin A.
- Continue gentle cleansing with water 12 hours after treatment. Avoid applying high pressure on skin when cleansing
- After treatment, avoid smoking, alcohol, sauna or excessive exercise.
- Drink 1.5 – 2 L of water daily.
- For post-treatment, application of regenerating moisturizing and anti-bacterial cream is recommended.
- Avoid applying high pressure (rubbing, scratching) on skin for 3-5 days after treatment.

Pink Clinical Result

Large pore & Irritated skin

3 treatments / 3weeks



Late 40s aged women

Dark circles

4 treatments / 4weeks



Early 30s aged man

Early 30s aged woman

Trouble & Irritated skin

5 treatments / 5weeks



Late 40s aged women

Trouble & Large pore

5 treatments / 5weeks



Early 30s aged women

Large pore

5 treatments / 5weeks





Pink

THE PINK-AGING TREATMENT

www.ribeskinx.com

Pink Single Package

